

Product Name:                    Contrac Soft Bait  
 Application No:                 86331/115500



Label Name:	Contrac Soft Bait
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	0.05 g/kg BROMADIOLONE Contains Denatonium Benzoate (Bittering Agent)
Mode of Action:	
Statement of Claims:	Highly palatable soft bait for the control of rats and mice including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.  Ready to use bait - Kills in a single feed. Soft Bait = palatable and attractive.
Net Contents:	Net 500 g Contains 50 X 10 g sachets which are illegal to sell separately.  Net 20 kg Contains 2000 X 10 g sachets which are illegal to sell separately.  DO NOT DESTROY CONTAINER WHILE PRODUCT STILL REMAINS.  10 g Not to be sold separately. Before use read all directions on the outer pack.
Restraints:	DO NOT exceed 3 m between bait stations for mice, or 9 m for rats. DO NOT place bait sachets in the open. DO NOT use bait stations further than 2 m from buildings.
Directions for Use:	DIRECTIONS FOR USE SITUATION In and around domestic homes, agricultural, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.

PEST  
Mice (*Mus musculus*)

Rats (*Rattus norvegicus*, *Rattus rattus*)

**RATE:**

Mice: 1-4 soft baits per bait station or placement.  
Rats: 2-20 soft baits per bait station or placement.

**CRITICAL COMMENTS**

Eliminate as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Place baits so as to prevent access to non-target animals. Tamper resistant bait stations must be used at locations where access to non-target animals cannot otherwise be prevented. Inspect bait stations on a regular basis. Replace eaten or old/deteriorated baits. If eaten quickly, increase the number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

Other Limitations:

Withholding Periods:

Trade Advice:

General Instructions:

**GENERAL INFORMATION**

Conrac Soft bait is a highly palatable soft bait for the control of rats and mice including those resistant to other anticoagulants. For use in and around buildings. One feed kills rats and mice. Single feed rodenticide.  
Soft Bait = palatable and attractive.

**GENERAL INSTRUCTIONS**

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoilt bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead animals and rodent faeces by burning or burial, and left over bait as per disposal instructions. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

Resistance Warning:

Precautions:

**PRECAUTION**

DO NOT remove bait from sachet.

The use of this product for the control of protected native rodents requires permission from wildlife authorities.

	<p>DO NOT place bait in places accessible to children, domestic animals livestock and non-target wildlife, unless in tamper resistant bait stations.</p> <p>DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed, or surfaces that come into direct contact with food. The product is restricted to use in an around buildings (within 2 metres) or enclosed spaces.</p> <p>DO NOT apply bait to crops.</p>
--	--

<p>Protections:</p>	<p><b>PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT</b></p> <p>DO NOT contaminate dams, rivers, streams or drains with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product and dead or affected rodents and rodent faeces are hazardous to pigs, cats, dogs, native wildlife and poultry.</p> <p><b>PROTECTION OF LIVESTOCK</b></p> <p>Bait, and dead or affected rodents are hazardous to birds, dogs, cats, pigs, poultry and non-target wildlife. Regularly search for, remove and destroy dead or affected rodents.</p>
---------------------	---

<p>Storage and Disposal:</p>	<p>Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away form children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. To the extent possible, untaken baits should be destroyed by burning or burial according to State or Territory requirements.</p>
------------------------------	---

<p>Safety Directions:</p>	<p>Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear rubber gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K (Phytomenadione) is antidotal.</p>
---------------------------	--

<p>First Aid Instructions:</p>	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Vitamin K (Phytomenadione) is antidotal.</p>
--------------------------------	--

<p>First Aid Warnings:</p>	
----------------------------	--