

MAIN PANEL

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Starycide[®]

INSECT GROWTH REGULATOR

ACTIVE CONSTITUENT: 48 g/L TRIFLUMURON

GROUP 15A INSECTICIDE

1 L

For the control silverfish, immature fleas and cockroaches in indoor and outdoor situations.

IMPORTANT:
READ THE ATTACHED BOOKLET BEFORE USE

Rear panel attached to container

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the nose, throat and skin. Avoid contact with skin. Do not inhale spray mist. When opening the container and preparing spray wear elbow length PVC gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 131126).

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval Number: 61251/1L/1207

Cislin, Tempo, Resposar and Starycide® are Registered Trademarks of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Website: www.bayeres.com.au

Batch Number:

DOM:

For 24 hour specialist advice in an emergency only phone 1800 033 111

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Starycide[®]
Insect Growth Regulator

ACTIVE CONSTITUENT: 48 g/L TRIFLUMURON

GROUP	15A	INSECTICIDE
--------------	------------	--------------------

For the control of silverfish, immature fleas and cockroaches in indoor and outdoor situations

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the nose, throat and skin. Avoid contact with skin. Do not inhale spray mist. When opening the container and preparing spray wear elbow length PVC gloves. Wash hands after use. After each day's use wash gloves.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 131126).

Directions for use:

SITUATIONS	RATE	PEST	CRITICAL COMMENTS
Domestic, commercial, industrial and public buildings: - including food processing establishments, shops, factories, ships, offices, schools, storerooms, hospitals, barracks, private houses and external areas associated with such establishments.	5 mL/L / 10m ²	Flea larvae	Thoroughly spray all infested areas. It is recommended that premises are vacuumed and contents of vacuum bag removed before treatment takes place. This is particularly important where adult fleas are present. Ensure pets are treated with a suitable registered insecticide to avoid re-infestation of the treated premises. Exterior Use: Prewetting the soil with a garden hose may be required on very dry soil.
	10 – 20 mL/ L water	Immature cockroaches	When treating for cockroaches apply thoroughly to cracks, crevices and other places where cockroaches may hide and breed. Apply to point of run-off. The lower (10 mL/L) rate will be effective in most situations for a full season. Use the higher rate under difficult conditions or where more than season long control is required.
		Silverfish	When treating for silverfish apply thoroughly to surfaces in cupboards, cracks, crevices and other places frequented by silverfish. Apply to point of run-off.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Starycide is a slow acting insect growth regulator which has its greatest effect on early juvenile stages of insects. Applied to breeding areas, Starycide will assist in preventing pest populations from building up to problem levels.

Where quick action is required the use of a knockdown product as a tank mix with Starycide is recommended. Starycide is compatible with most adulticides such as Cislin, Tempo and Responsar. As an insect growth regulator, without the excito-repellent action of a synthetic pyrethroid Starycide will also complement treatment with gel baits for cockroach control.

If used alone on established pest populations, it will take several weeks for populations to decline to acceptable levels (3-4 weeks for fleas and 1-2 months for cockroaches). Starycide should only be used alone when the pest population has already been reduced by other measures or when the property owner is willing to accept a slow action.

Mixing:

Shake product well before use. Mix Starycide in the required amount of water or prepared knockdown product to uniformly and accurately apply the required amount.

Application

Do not spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

Inside booklet contd.

GENERAL PRECAUTIONS:

- Before use - remove or cover all exposed foodstuffs; cover all dishes and utensils, and places where food is prepared or stored.
- After use - thoroughly ventilate treated area. Clean up thoroughly before processing/serving resumes.
- When using in animal houses do not spray directly onto livestock, their food, water, litter, feed bins, or surfaces that are accessible to the stock.
- Do not spray directly on humans, exposed food or food utensils or food preparation areas.
- Avoid contact of spray with timber to which a stain is subsequently to be applied, unless the owner is notified that the timber will need additional preparation, consistent with that if water alone had contacted the timber. (Evidence of water contact on timber is sometimes only evident once the stain has been applied).

INSECTICIDE RESISTANCE WARNING

GROUP	15A	INSECTICIDE
--------------	------------	--------------------

For insecticide resistance management Starycide is a Group 15A insecticide. Some naturally occurring insect biotypes resistant to Starycide and other Group 15A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Starycide or other Group 15A insecticides are used repeatedly. The effectiveness of Starycide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from failure of Starycide to control resistant insects.

For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to non-target insects, fish, yabbies and other aquatic and soil arthropods. Do NOT contaminate dams, drains, ponds, creeks and other waterways with the product or used containers.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval Number: 61251/1207

Cislin, Tempo, Responsar and Starycide® are Registered Trademarks of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Website: www.bayeres.com.au

Batch Number:

DOM:

For 24 hour specialist advice in an emergency only phone 1800 033 111
